

GROUP 25 FUNGICIDE

Agri-Mycin® 50

Agricultural Streptomycin

ACTIVE INGREDIENT:

Streptomycin Sulfate (Equivalent to 50% streptomycin base)..... 65.8%

OTHER INGREDIENTS: 34.2%

TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION / PRECAUCIÓN
 Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

For Medical Emergencies Only, Call (877) 325-1840

EPA Reg. No. 55146-98

Net Weight

Manufactured for
 Nufarm Americas Inc.
 AGT Division
 11901 S. Austin Avenue
 Alsip, IL 60803

EPA Est. No. 67545-AZ-001

3.0 Pounds



FIRST AID	
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION / PRECAUCIÓN

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear
- A NIOSH approved particulate respirator with an N, R, or P filter, NIOSH approval number prefix TC-84A.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS	
Users should:	<ul style="list-style-type: none"> Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product may be hazardous to aquatic plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This material is not to be used for medical, veterinary, or human purposes.

RESISTANCE MANAGEMENT

This product contains a Group 25 fungicide/bactericide and is effective against pathogens which are resistant to fungicides/bactericides with modes of action different from those of fungicide Group 25. Pathogen biotypes resistant to streptomycin may eventually dominate the fungal/bacterial population if Group 25 fungicides/bactericides are used repeatedly in the same site or in successive years as the primary method of control for targeted pathogen species. Selection for resistance may be particularly rapid if resistance to Group 25 fungicides/bactericides is already present in the pathogen population. Making more than two consecutive applications of this product may be conducive to the selection of resistant strains.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried

CROP USE DIRECTIONS
MIXING INSTRUCTIONS

Dilution Table

Concentration Desired (ppm)*	Quantity of Product per Volume of Water				
	5 gal	50 gal	100 gal	500 gal	600 gal
50	1/2 tsp	2/3 oz	1 1/3 oz	6 2/3 oz	8 oz
60	2/3 tsp	0.8 oz	1.6 oz	8 oz	9.6 oz
100	1 tsp	1 1/3 oz	2 2/3 oz	13 1/3 oz	1 lb
200	2 tsp	2 2/3 oz	5 1/3 oz	1.7 lb	2 lb

* Parts per million of streptomycin base in prepared solution

ROW CROPS

CROP	DISEASE	PRODUCT RATE OR CONCENTRATION	USE INSTRUCTIONS
POTATO	Soft Rot Blackleg	100 ppm (Refer to Dilution Table above)	Soak cut seed pieces in solution for several minutes and plant as usual. <i>Note:</i> A suitable fungicide (such as Maxim® PSP, Maxim MZ Fungicide, Captan, or a product containing a dithiocarbamate a.i.) should be used as an adjunct to this treatment for the control of fungal diseases associated with potato seed pieces.
		200 ppm (equivalent to 2 2/3 - 8 oz in 100 - 300 gal/A)	
TOBACCO	Wildfire Blue Mold	<i>For preventive action</i> 100 ppm (equivalent to 5 1/3 oz in 200 gal/A or 25 gal/5,000 ft²)	Apply first spray when plants are in the 2-leaf stage or about the size of a dime or when blue mold first appears in the area. Continue applications at 5- to 7-day intervals until plants are set in the field. Additional protection may be obtained by spraying field plants weekly.
		<i>For curative action</i> 200 ppm (equivalent to 10 2/3 oz in 200 gal/A)	
		200 ppm (equivalent to 1 1/3 - 5 1/3 oz in 25 - 100 gal/A)	Apply in locations where wildfire has been a problem in recent years or where applications have been delayed until disease appears. Follow the same schedule as above.

TREE CROPS

CROP	DISEASE	PRODUCT RATE OR CONCENTRATION	USE INSTRUCTIONS
APPLE	Fire Blight	8 - 16 oz/A (equivalent to 50 - 100 ppm at 600 gal/A)	Make first application at 20 - 30% bloom. Spray trees every 3 to 4 days during blossom time. After petal fall, continue applications at 10- to 14-day intervals to control twig blight (this could mean an additional 6 to 8 applications after blossom sprays).
		Restriction Do not apply within 50 days of harvest.	
PEAR (Except CA)	Fire Blight	8 - 16 oz/A (equivalent to 50 - 100 ppm at 600 gal/A)	Make first application at 20 - 30% bloom. Spray trees every 3 to 4 days during blossom time. After petal fall, continue applications at 10- to 14-day intervals to control twig blight (this could mean an additional 6 to 8 applications after blossom sprays).
		Restriction Do not apply within 30 days of harvest.	
PEAR (California only)	Fire Blight	9.6 oz/A (equivalent to 60 ppm at 600 gal/A)	Make first application at 10% bloom and repeat at 5-day intervals until all late bloom is over. Continue applications at 5- to 7-day intervals to control shoot and fruit infections.
		Restriction Do not apply within 30 days of harvest.	

VEGETABLE CROPS

(Seedlings of the below listed crops grown in greenhouses and similar structures)

CROP	DISEASE	PRODUCT RATE OR CONCENTRATION	USE INSTRUCTIONS
CELERY (Florida only)	Bacterial Blight	200 ppm (equivalent to 2 2/3 - 5 1/3 oz in 50 - 100 gal/A)	Apply first spray when seedlings are in the 2-leaf stage, when first true leaves appear. Continue applications at 4- to 5-day intervals until transplanted in the field.
PEPPER	Bacterial Spot Bacterial Speck*	200 ppm (equivalent to 2 2/3 - 5 1/3 oz in 50 - 100 gal/A)	Apply first spray when seedlings are in the 2-leaf stage, when first true leaves appear. Continue applications at 4- to 5-day intervals until transplanted in the field.
TOMATO	Bacterial Spot Bacterial Speck*	200 ppm (equivalent to 2 2/3 - 5 1/3 oz in 50 - 100 gal/A)	Apply first spray when seedlings are in the 2-leaf stage, when first true leaves appear. Continue applications at 4- to 5-day intervals until transplanted in the field.

*Not registered for use in California

ORNAMENTAL PLANTS AND SHRUBS

Grown in Greenhouses, Shadehouses, Fields, Nurseries (container, bench, or bed-grown), and Residential & Commercial Landscapes

*Intended for use by professional applicators

PLANT	DISEASE	PRODUCT RATE OR CONCENTRATION	USE INSTRUCTIONS
Garden: Apple trees Pear trees Ornamentals: Amelanchier* Cotoneaster* Pyracantha	Fire Blight	100 ppm (equivalent to 2 2/3 - 8 oz in 100 - 300 gal/A)	Apply in foliar and blossom sprays. Apply first spray at start of blossoming period. Continue applications every 3 to 4 days during blossom time. Apply additional sprays every 5 to 7 days after blossom period when weather favors spread of fire blight.
		Restriction Do not apply after fruit is visible.	
Berberis* Carnation* Forsythia* Geranium* Hedera (ivy)* Hydrangea* Impatiens* Lonicera (honeysuckle)* Philadelphus* Philodendron Poinsettia* Rudbeckia (Black-eyed susan)* Salvia* Syringa* Viburnum*	Bacterial Leaf Spot* Bacterial Leaf Rot Bacterial Blight* (<i>Pseudomonas</i> spp., <i>Xanthomonas</i> spp.)	200 ppm (equivalent to 5 1/3 oz - 1 lb in 100 - 300 gal/A)	Apply as preventive spray or at first sign of water-soaked areas on leaves. Repeat at 4- to 5-day intervals. <i>For curative action</i> Remove all rotted leaves from plant and then spray every 4 days.
Chrysanthemum Hydrangea*	Bacterial Wilt (<i>Pseudomonas solanacearum</i>)	50 ppm (Refer to Dilution Table above)	Soak plant cuttings in solution for 4 hours. Plant as usual.
Calla Lily Gladiolus* Orchid*	Leaf Spot* Soft Rot caused by <i>Erwinia</i> spp.	100 - 200 ppm (Refer to Dilution Table above)	<i>Pre-Plant Treatment</i> Apply as a postharvest, pre-plant dip or spray onto exposed bulbs.
		100 - 200 ppm (equivalent to 2 2/3 - 5 1/3 oz per 100 gal)	<i>Post-Plant, Pre-Emergent Drench</i> Apply as a drench at a rate of 3 to 4 ounces of solution per 6 inch pot or 3 to 4 ounces of solution per setting in the field.
		200 ppm (equivalent to 5 1/3 oz - 1 lb in 100 - 300 gal/A)	<i>Preventive treatment*</i> Apply as a foliar spray at first sign of water-soaked areas on leaves. Continue applications at 4- to 7-day intervals.

(continued)

50"

ACCEPTED
FOR REGISTRATION

09/28/2017

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

Doc ID 553608

Classified for
"RESTRICTED USE"
in New York State
under 6NYCRR Part 326

RV052316