LEBANON	ACCEPTED VIA NOTIFICATION LABEL NOT REVIEWED
TREFLAN* 5G CONTAINS TREFLAN* HERBICIDE ACTIVE INGREDIENT:	Apr 28, 2011 New York State Department of Environmental Conservation Division of Materials Management Pesticide Product Registration
trifluralin OTHER INGREDIENTS: TOTAL. *Trademark of Dow AgroSciences LLC. (L) (D) EPA Reg. No. 961-405 Contains 2 pounds active ingredient per 40 pound bag. EPA Est. No. 961-PA-01: 961-IL-01 Superscript used is first letter of lot number. Manufactured using the processes of U.S. Patents 4,120,905 and 4,226,789. 24-366883	

Doc Id: 526383

LEBANON TREFLAN* 5G is a selective preemergence herbicide for control of certain annual grasses and broadleaf weeds in:

- Container Grown Ornamentals
- Landscape Ornamentals
- Nursery Stock
- Ground Covers
- Non-Cropland

- Ornamental Bulbs
- Non-Bearing Fruit and Nut Trees and
 - Non-Bearing Vineyards in Ornamental Landscaping
- Established Flowers

KEEP OUT OF REACH OF CHILDREN CAUTION

In case of an emergency endangering life or property involving this product, call Chemtrec at 1-800-424-9300.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

	FIRST AID
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
	Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice.
	Have person sip a glass of water if able to swallow.
	Do not induce vomiting unless told to by a poison control center or doctor.
	Do not give anything to an unconscious person.
IF ON SKIN:	Take off contaminated clothing.
	Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air.
	 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
	Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For Emergency Medical Information call (888) 208-1368.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other Handlers must wear:

- · Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- . Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- · Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and ovster. Do not apply in a manner which will directly expose canals. lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

NOT FOR USE ON PLANTS BEING GROWN FOR SALE OR OTHER COMMERCIAL USE, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES. FOR USE ON PLANTS INTENDED FOR AESTHETIC PURPOSES OR CLIMATIC MODIFICATIONS AND BEING GROWN IN INTERIOR PLANTSCAPES, ORNAMENTAL GARDENS OR PARKS, OR ON GOLF COURSES OR LAWNS AND GROUNDS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

ENTRY RESTRICTIONS

Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is drv after the watering-in.

OFNEDAL INFORMATION

ury aller the watering-in.

GENERAL INFORMATION

This product is a preemergence herbicide for control of certain annual grasses and broadleaf weeds in container grown ornamentals, landscape ornamentals, nursery stock, ground covers, established flowers, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards in ornamental landscaping only, and noncropland. Apply before or after planting but prior to germination of target weeds, or immediately after cultivation. Length of weed control will vary with weed population, potting media or soil conditions, temperature, watering regime and other factors. Following application, user should monitor and observe level of weed control over time to determine when additional applications may be needed. Repeat applications should not be made sooner than 60 days after a previous application. Do not apply over 240 lbs. per acre total of this product within a 12-month period.

GENERAL USE PRECAUTIONS

This product does not control established weeds. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings and trash should be removed or thoroughly mixed into soil prior to treatment. Soil should be in good condition and free of clods at the time of application. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation, is required to activate this product. Optimum weed control is obtained when this product is activated within 3 days of application. If rainfall or irrigation has not occurred within 3 days of application and tillage is possible, this product may be activated using cultivation equipment capable of uniformly mixing the herbicide into the upper 1-2 inches of soil. Failure to activate this product within 3 days of application may result in erratic weed control. Do not apply when wind conditions favor drift of the granules from the target area.

Optimum weed control will be obtained when followed by overhead irrigation or rainfall within a few hours after surface application.

WEEDS CONTROLLED WHEN APPLIED AT 80 LBS./ACRE (1.8 Lbs./1,000 Sq. Ft.)

		ANNUA	AL GRASSES					
Barley, hare	Brome	Cupgrass, Southwestern	Goosegrass	Oat, Wild	Signalgrass			
Barnyardgrass (watergrass)	Cheat	Fescue, rattail	Junglerice	Panicum, fall	Sprangletop, Mexican			
Bluegrass, annual	Crabgrass	Foxtail	Lovegrass	Sandbur, field	Stinkgrass			
					Witchgrass			
	ANNUAL BROADLEAF WEEDS							
Chickweed, common	Knotweed, prostrate	Lambsquarters, common	Pigweed	Purslane, common	Thistle, Russian			
Goosefoot, nettleleaf	Kochia	Nettle, stinging	-					
			l -+ 00 lb / (4 0 lb /400	0 #)				
In addition to the weeds controlled,	In addition to the weeds controlled, the following weeds will be partially controlled or suppressed at 80 lbs./acre (1.8 lbs./1000 sq. ft.).							
Carpetweed	Johnsongrass (from seed)	Panicum, Texas	Pusley, Florida	Shattercane				

LebanonTurf

For technical assistance or more information about our products visit www.LebanonTurf.com

Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

Made in the U.S.A. www.LebanonTurf.com

NET WT. 40 LBS. (18.14 kg)

APPLICATION DIRECTIONS

Apply this product using a drop or rotary-type spreader designed to apply granular herbicides or insecticides. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or plant injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

APPLICATION TECHNIQUES

• When using a drop-type spreader, a splash board mounted under the hopper will provide more even granule distribution.

• A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.

• For treating smaller areas or rows of nursery stock or ornamental beds, a hand held or push-type rotary applicator such as whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed. A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, hard to treat areas. Carefully measure the amount of product needed to avoid over-application.

APPROVED USES

Container Grown Ornamentals, Landscape Ornamentals, Nursery Stock, Ground Covers, Established Flowers, Ornamental Bulbs, Non-bearing Fruit and Nut Trees, Non-bearing Vinevards in Ornamental Landscaping Only, and Non-cropland. This product is recommended as a preemergence treatment for control of certain annual grasses and broadleaf weeds in container grown ornamentals, landscape ornamentals, nursery stock, ground covers, established flowers, ornamental bulbs, non-bearing fruit and nut trees, non-bearing vineyards in ornamental landscaping only, and non-cropland. Apply before or after planting but prior to germination of target weeds, or immediately after cultivation.

Refer to the "General Information" section of this label for general use precautions and information on application, application rates and weeds controlled.

SPECIAL USE PRECAUTIONS

To avoid possible injury, do not apply this product to:

. Unrooted liners or cuttings that have been planted in pots for the first time

· Pots less than four inches wide

· Ground covers until they are established and well rooted

Do not apply to newly transplanted ornamentals, nursery stock, ground covers, flowers and non-bearing fruit and nut crops and non-bearing vineyards in ornamental landscaping until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or injury may occur.

Do not make preplant applications of this product to areas where gladioli corms are less than one inch in diameter will be planted or injury may occur.

Do not apply this product in greenhouse or other enclosed structures.

Carob

Redbud

Repeat applications should not be made sconer than 60 days after a previous application of this product. Do not apply over 240 lbs, per acre total of this product within a 12 month period.

Prior to using this product on a large number of plants, the treated plants should be observed for signs of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. THE USER ASSUMES RESPONSIBILITY FOR ANY PLANT DAMAGE OR OTHER LIABILITY RESULTING FROM THE USE OF THIS PRODUCT ON PLANT SPECIES NOT RECOMMENDED ON THIS LABEL.

This product may be used on the following established plant species when container grown or field grown:

Balsam Fir White Fir Albus Flowering Maple Luteus Flowering Maple Roseus Flowering Maple Tangerine Flowering Maple Vesuvius Red-Flowering Maple Flame Maple Norway Maple Red Maple Red Sunset Maple Silver Maple Sugar Maple Queen Palm River Birch Paper Birch

European White Birch Kousa Dogwood Bottle Tree Green Hawthorn Black Olive Carrot Wood Chinese Chestnut Arizona Cypress Russian Olive Redoum Eucalvotus Filicoides-Fernspray Cypress Mealy Eucalyptus Gracilis-Slender Hinoki Cypress Silver Dollar Eucalyptus Swara False Cypress Coolibah Tree Red Ironbark Eucalyptus Squarrosa-Moss Cypress Palm (Chamaedorea cataractarum) Ficus Palm (Chamaedorea costaricana) Mini Fucus Parlor Palm White Ash Cloud Nine Dogwood Shamel Ash Flowering Dogwood Ginko-Maidenhair Tree

TREES

Honey Locust Shademaster Honey Locust Tovon Florida Anise-Tree Eastern Red Cedar Japanese Larch American Sweet Gum Tuliptree Southern Magnolia Crabapple White Mulberry Banana Blackoum Sourwood Norway Spruce

Pendula-Weeping Norway Spruce Repens-Spreading Norway Spruce Dwarf Alberta Spruce White spruce Dwarf Globe Blue Spruce Glauca-Colorado Blue Spruce Hoopsii-Hoop's Blue Spruce Koster-Koster Blue Spruce Bristlecone Pine Canary Island Pine Shore Pine, Beach Pine Eldarica Pine Bosnian Pine Pumilio-Shrubby Swiss Mountain Pine Austrian Black Pine

Monterev Pine Red Pine White Pine Columnar Scotch Pine Scotch Pine Loblolly Pine Japanese Black Pine London Planetree American Sycamore California Sycamore Podocarpus Cottonwood Chilean Mesquite Yoshino Flowering Cherry Douglasfir

Bear Oak Pin Oak Willow Oak Red Oak Live Oak Black Locust Willow Giant Seguoia Mahogany Yellow Tab Baldcypress Eastern Hemlock Chinese Elm Mexican Fan Palm

Scarlet Oak

ORNAMENTAL SHRUBS

Paper Birch

Edward Goucher Abelia Glossy Abelia Abyssinica Acacia Prostrate Acacia Shoestring acacia Copper Leaf Amur Maple Coral Bark Japanese Maple Dwarf Japanese Maple Century Plant False Spirea Japanese Painted Fern Covotebush William Penn Barberry Mentor Barberry Atropurea-Redleaf Japanese Barberry Aurea-Golden Japanese Barberry Crimson Pygmy Barberry Rose Glow Barberry Barbara Karst California Gold Pink Pixie Scarlet O'Hara Temple Fire Texas Dawn

Wooly yarrow Lily-of-the-Nile Beecharass Cape Weed Thrift Asparagus Fern Bellflower Variegated Carex

Yarrow Floss Flower Alyssum Snapdragon African Daisy Dusty Miller Aster (perennial) Calendula

Japanese Boxwood Korean Boxwood Common Boxwood Lemon Bottlebrush Weeping Bottlebrush Spring Torch Scotch Heather Japanese Camellia Sasangua Camellia Feathery Cassia Wild Lilac Plum Yew Snow-in-Summer Kosteri Cypress Nana-Dwarf Hinoki Cypress Torulosa Cypress Filifera-Thread Cypress Areca palm Summersweet Japanese Cleyera Sibirica-Siberian Dogwood Bailevi Red Osier Dogwood Flaviramea-Yellowtwig Dogwood Royal Purple Smoke Tree Coral Beauty Smoke Tree Eichholz Smoke Tree

Largeleaf Iceplant

Dwarf Plumbago

Rockrose

Coreopsis

Crown Vetch

Cotoneaster

Cornflower

Sweet Sultan

Morningglory

Coreopsis

Cosmos

Dahlia

Velvet Centaurea

Chrvsanthemum

White Iceplant

Pampas Grass

Descampsia Trailing Rosea Iceplant Iceplant Blue Fescue Strawberry, Beach Gazania Golden Hakonechloa Algerian Ivy

Praecox-Early Cotoneaster

Cranberry Cotoneaster

Pyrenees Cotoneaster

Bearberry Cotoneaster

Himalayan Cotoneaster

Hollandia-Warminster Broom

Rock Cotoneaster

Zabel Cotoneaster

Lena-Scotch Broom

Fruitland Silver Berry

Mediterranean Pink Heather

Canadale Gold Euonymus

Emerald'n Gold Euonymus

Wintercreeper Euonymus

Variegated Evergreen Euonymus

Silver King-Euonymus

Spreading Euonymus

Purple Bell Heather

Cornish Heather

Winged Euonymus

Sunspot Euonymus

Fragrant Daphne

Hopseed Bush

Sago Palm

Deutzia

Dianthus

Gaillardia Geum Gladiolus

ORNAMENTAL SHRUBS

Pineapple Guava Forsythia August Beauty Gardenia Gardenia Radican Gardenia Salal/Lemon Leaf Carolina jessamine Woadwaxen Ross Estev-Hibiscus Rose-of-Sharon- Heart Rose-of-Sharon- Red Bird Rose-of-Sharon-Woodbridge Holly Mystery Gardenia Henry Garnet Holly Leaf Sweetspire Ixora Juniper Mountain Laurel Crape Myrtle Lantana Coast Leucothoe Drooping leucothoe Privet Chinese Fountain Palm Flowering Woodbine

GROUND COVERS

English Ivv Daylilly Rupturewort Albo-Marginata Hosta Aaronsbeard St. Johnswort Angelwing Jasmine Trailing Iceplant

ESTABLISHED FLOWERS

Sunflower Petunia Pincushion Flower Sweet Alyssum Marigold, Cape Balsam Lupine Phlox Lamb's Ears Snow on the Mountain Impatiens Stock Portulaca Stoke's Aster Ixora Four O'Clock Rose Marigold Sweet Pea Forget-Me-Not Nasturtium Blackeyed Susan Statice Nicotiana Golden Glow Vinca Zinnia Baby's Breath Lobelia Poppy, California Salvia

Serotina Woodbine

Trumpet Honevsuckle

Leather Leaf Mahonia

Compacta-Dwarf Heavenly Bamboo

Harbour Dwarf-Heavenly Bamboo

Nana Compacta-Heavenly Bamboo

Nana Purpurea-Heavenly Bamboo

Woods Dwarf Heavenly Bamboo

Creeping Mahonia

Heavenly Bamboo

Hardy Red Oleander

Ruby Lace Oleander

Fortunes Osmanthus

Japanese Andromeda

Mountain Fire Lily-of-the-Valley

Snowdrift Lily-of-the-Valley

Templebells Lily-of-the-Valley

Valley Rose Lily-of-the-Valley

Valley Valentine Lily-of-the-Valley

Forest Flame Lilv-of-the-Valley

Pigmy Date Palm

Fraser's Photinia

White Lilv Turf

Big Blue Lily Turf

Magestic Lily Turf

Eulalia Grass

Lilac Beauty Lilv Turf

Silvery Sunproof Lilv Turf

Variegated Lirope Lilv Turf

Green/Creeping Lily Turf

Oleander

Mockorange

Wax Myrtle

ORNAMENTAL BULBS

This product may be applied for control of susceptible annual weeds in ornamental bulbs, e.g., bulbous iris, daffodil (narcissus), hyacinth and tulip. Apply to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. This product may also be applied following bulb emergence. For fall planted bulbs, apply this product again in later winter or early spring to weed-free soil surfaces.

NON-BEARING FRUIT AND NUT TREES AND NON-BEARING VINEYARDS IN ORNAMENTAL LANDSCAPING ONLY*

Almond	Boysenberry	Fig	Kiwi	Orange	Pomegranate
Apple	Cherry, sour	Filbert	Lemon	Peach	Prune
Apricot	Cherry, sweet	Gooseberry	Loganberry	Pear	Raspberry
Avocado	Currant	Grape, American	Macadamia Nut	Pecan	Walnut, black
Blackberry	Dewberry	Grape, European	Nectarine	Pistachio	Walnut, English
Blueberry	Elderberry	Grapefruit	Olive	Plum	-

*Non-bearing fruit and nut trees and non-bearing vinevards are defined as plants which will not bear fruit for at least one year after treatment.

Muao-Muaho Pine Green Pittosporum Japanese Pittosporum Wheeler's Dwarf Pittosporum Blue Cape Plumbago Plumbago Yewpine Tassel Fern Cinquefoil Carolina Laurel Cherry Dwarf Pink Flowering Almond Pyracantha Charisma-Monruce Rhaphiolepis Enchantress-Moness Rhaphiolepis India Hawthorn Springtime-Monme Rhaphiolepis Roundleaf Rhaphiolepis Azalea/Rhododendron Africa Sumac Ramanas Rose Rosemarv Japanese Skimmia Reeve's Skimmia Anthony Waterer Spiraea Dolchia Spiraea

Creeping Wirevine

Dwarf Mondo Grass

Trailing African Daisy

Japanese Spurge

Fountain Grass

Myoporum

Mondo Grass

Japanese Alpine Spiraea Shirobana Spiraea Bridal Wreath Chinese Lilac Common Lilac Upright Japanese Yew Anglojap Yew Cape Honeysuckle Ternstroemia gymnanthera American Arborvitae Emerald Arborvitae Globosa-Globe Arborvitae Little Giant-Dwarf Arborvitae Nigra-Dark American Arborvitae Pyramidalis-Pyramid Arborvitae Rheingold Arborvitae Techny Arborvitae Aureus Nana-Dwarf Golden Arborvitae Minima Glauca-Dwarf Arborvitae Christmas Palm Virburnum Weigela **Xylosma** Yucca

Ribbon Grass Stonecrop (Sedum) Germander Asaian Jasmine Verbena Speedwell Periwinkle

NON-CROPLAND AREAS

This product is recommended as a preemergence herbicide for control of certain annual grasses and broadleaf weeds on industrial sites, utility substations, highway guard rails, sign posts and delineators.

Apply anytime prior to germination of target weeds. Areas to be treated should be free of established weeds or existing weeds should be controlled with postemergence herbicides.

Refer to the "General Information" section prior to using this product on non-cropland.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store product in its original container in a cool, dry, locked place out of the reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Lebanon Seaboard Corporation. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, LEBANON SEABOARD CORPORATION MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Lebanon Seaboard Corporation is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Lebanon Seaboard Corporation DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT LEBANON SEABOARD CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.

LebanonTurf

For technical assistance or more information about our products visit www.LebanonTurf.com

Manufactured by: Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

Made in the U.S.A. www.LebanonTurf.com

NET WT. 40 LBS. (18.14 kg)



Lebanon TREFLAN* 5G is a selective preemergence herbicide for control of certain annual grasses and broadleaf weeds in:

- Container Grown Ornamentals
 Ground Covers
- Landscape Ornamentals
- Established Flowers
- Non-Bearing Fruit and Nut Trees and Non-Bearing Vineyards in Ornamental Landscaping



• Ornamental Bulbs Noncropland

ACTIVE INGREDIENT:

trifluralin OTHER INGREDIENTS:	
Contains 2 pounds active ingredient per 40 pound bag.	(L) (D)
EPA Reg. No. 961-405	EPA Est. No. 961-PA-01:961-IL-01
Trademark of Dow AgroSciences LLC	Superscript used is first letter of lot number

rademark of Dow AgroSciences LLC Manufactured using the processes of U.S. Patents 4,120,905 and 4,226,789.

KEEP OUT OF REACH OF CHILDREN CAUTION

Refer to back panel of bag for precautionary information and Directions for Use including STORAGE AND DISPOSAL.

In case of an emergency endangering life or property involving this product, call Chemtrec at 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

FIRST AID

	IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
I		Call a poison control center or doctor for treatment advice.
I	IF SWALLOWED:	 Call a poison control center or doctor immediately for treatment advice.
I		Have person sip a glass of water if able to swallow.
I		Do not induce vomiting unless told to by a poison control center or doctor.
I		Do not give anything to an unconscious person.
I	IF ON SKIN:	Take off contaminated clothing.
I		Rinse skin immediately with plenty of water for 15-20 minutes.
I		Call a poison control center or doctor for treatment advice.
I	IF INHALED:	Move person to fresh air.
I		 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
I		Call a poison control center or doctor for further treatment advice.
	Have the product co	ntainer or label with you when calling a poison control center or doctor or going for treatment. For Emergency Medical Information call (888) 208-1368.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other Handlers must wear:

· Long-sleeved shirt and long pants

Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

- · Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - · Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.



For technical assistance or more information about our products visit www.LebanonTurf.com Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

Made in the U.S.A. www.LebanonTurf.com

NET WT. 40 LBS. (18.15 KG)

Lebanon TREFLAN* 5G is a selective preemergence herbicide for control of certain annual grasses and broadleaf weeds in:

- Container Grown Ornamentals
 Ground Covers
- Landscape Ornamentals
- Nursery Stock

- Established Flowers
- Ornamental Bulbs
- Non-Bearing Fruit and Nut Trees and Non-Bearing
- Vineyards in Ornamental Landscaping
- Noncropland

ACTIVE INGREDIENT:

trifluralin	5.0%
OTHER INGREDIENTS:	
TOTAL ·····	100.0%
Contains 2 pounds active ingredient per 40 pound bag.	(L) (D)
EPA Reg. No. 961-405	(L) (D) EPA Est. No. 961-PA-01:961-IL-01
Trademark of Dow AgroSciences LLC.	Superscript used is first letter of lot number.

Manufactured using the processes of U.S. Patents 4,120,905 and 4,226,789.

KEEP OUT OF REACH OF CHILDREN CAUTION

Refer to back panel of bag for precautionary information and Directions for Use including STORAGE AND DISPOSAL.

In case of an emergency endangering life or property involving this product, call Chemtrec at 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

	FIRST AID						
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a price control con						
IF SWALLOWED:	Call a poison control center or doctor for treatment advice. Call a poison control center or doctor immediately for treatment advice.						
	• Have person sip a glass of water if able to swallow.						
	Do not induce vomiting unless told to by a poison control center or doctor.						
IF ON SKIN:	Do not give anything to an unconscious person. Take off contaminated clothing.						
IF ON SKIN.	Rinse skin immediately with plenty of water for 15-20 minutes.						
	Call a poison control center or doctor for treatment advice.						
IF INHALED:	Move person to fresh air.						
	If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.						
	Call a poison control center or doctor for further treatment advice.						
Have the product col	Have the product container or label with you when calling a poison control center or doctor or going for treatment. For Emergency Medical Information call (888) 208-1368.						
	PERSONAL PROTECTIVE EQUIPMENT (PPE):						

Applicators and other Handlers must wear:

· Long-sleeved shirt and long pants

Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.



For technical assistance or more information about our products visit www.LebanonTurf.com

Manufactured by: Lebanon Seaboard Corporation 1600 E. Cumberland St. • Lebanon, PA 17042 Made in the U.S.A.

www.LebanonTurf.com

NET WT. 40 LBS. (18.15 KG)





DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

NOT FOR USE ON PLANTS BEING GROWN FOR SALE OR OTHER COMMERCIAL USE, OR FOR COMMERCIAL SEED PRODUCTION OR FOR RESEARCH PURPOSES. FOR USE ON PLANTS INTENDED FOR AESTHETIC PURPOSES OR CLIMATIC MODIFICATIONS AND BEING GROWN IN INTERIOR PLANTSCAPES, ORNAMENTAL GARDENS OR PARKS, OR ON GOLF COURSES OR LAWNS AND GROUNDS. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

ENTRY RESTRICTIONS

Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

GENERAL INFORMATION

This product is a preemergence herbicide for control of certain annual grasses and broadleaf weeds in container grown ornamentals, landscape ornamentals, nursery stock, ground covers, established flowers, ornamental bulbs, non-bearing fruit and nut trees, non-bearing vineyards in ornamental landscaping only, and noncropland. Apply before or after planting but prior to germination of target weeds, or immediately after cultivation. Length of weed control will vary with weed population, potting media or soil conditions, temperature, watering regime and other factors. Following application, user should monitor and observe level of weed control over time to determine when additional applications may be needed. Repeat applications should not be made sooner than 60 days after a previous application. Do not apply over 240 lbs. per acre total of this product within a 12-month period.

GENERAL USE PRECAUTIONS

This product does not control established weeds. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings and trash should be removed or thoroughly mixed into soil prior to treatment. Soil should be in good condition and free of clods at the time of application. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation, is required to activate this product. Optimum weed control is obtained when this product is activated within 3 days of application has not occurred within 3 days of application and tillage is possible, this product may be activated using cultivation equipment capable of uniformly mixing the herbicide into the upper 1-2 inches of soil. Failure to activate this product within 3 days of application may result in erratic weed control. Do not apply when wind conditions favor drift of the granules from the target area.

Optimum weed control will be obtained when followed by overhead irrigation or rainfall within a few hours after surface application.

WEEDS CONTROLLED WHEN APPLIED AT 80 LBS./ACRE (1.8 Lbs./1,000 Sq. Ft.)

ANNUAL GRASSES								
Barley, hare	Brome	Crabgrass	Fescue, rattail	Goosegrass	Lovegrass	Panicum, fall	Signalgrass	Stinkgrass
Barnyardgrass (watergrass)	Cheat	Cupgrass, Southwestern	Foxtail	Junglerice	Oat, Wild	Sandbur, field	Sprangletop, Mexican	Witchgrass
Bluegrass, annual								
			ANNU	AL BROADLEAF WEEDS				
Chickweed, common	Goosefoot, nettleleaf	Knotweed, prostrate	Kochia	Lambsquarters, common	Nettle, stinging	Pigweed	Purslane, common	Thistle, Russian
In addition to the weeds cor Carpetweed	ntrolled, the following weeds Johnsongrass (from seed)	s will be partially controlled or s Panicum, Texas	uppressed at 80 lbs. Pusley, Florida	/acre (1.8 lbs./1000 sq. ft.). Shattercane				
•	,							

APPLICATION DIRECTIONS

Apply this product using a drop or rotary-type spreader designed to apply granular herbicides or insecticides. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or plant injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

APPLICATION TECHNIQUES

• When using a drop-type spreader, a splash board mounted under the hopper will provide more even granule distribution.

- A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.
- For treating smaller areas or rows of nursery stock or ornamental beds, a hand held or push-type rotary applicator such as whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.

• A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, hard to treat areas. Carefully measure the amount of product needed to avoid over-application.

APPROVED USES

Container Grown Ornamentals, Landscape Ornamentals, Nursery Stock, Ground Covers, Established Flowers, Ornamental Bulbs, Non-bearing Fruit and Nut Trees, Non-bearing Vineyards in Ornamental Landscaping Only, and Non-cropland.

This product is recommended as a preeemergence treatment for control of certain annual grasses and broadleaf weeds in container grown ornamentals, landscape ornamentals, nursery stock, ground covers, established flowers, ornamental bulbs, non-bearing fruit and nut trees, non-bearing vineyards in ornamental landscaping only, and non-cropland. Apply before or after planting but prior to germination of target weeds, or immediately after cultivation.

Refer to the "General Information" section of this label for general use precautions and information on application, application rates and weeds controlled.

SPECIAL USE PRECAUTIONS

To avoid possible injury, do not apply this product to:

• Unrooted liners or cuttings that have been planted in pots for the first time

• Pots less than four inches wide

Ground covers until they are established and well rooted

Do not apply to newly transplanted ornamentals, nursery stock, ground covers, flowers and non-bearing fruit and nut crops and non-bearing vineyards in ornamental landscaping until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or injury may occur.

Do not make preplant applications of this product to areas where gladioli corms are less than one inch in diameter will be planted or injury may occur.

Do not apply this product in greenhouse or other enclosed structures.

Repeat applications should not be made sooner than 60 days after a previous application of this product. Do not apply over 240 lbs. per acre total of this product within a 12 month period.

Prior to using this product on a large number of plants, the treated plants should be observed for signs of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. THE USER ASSUMES RESPONSIBILITY FOR ANY PLANT DAMAGE OR OTHER LIABILITY RESULTING FROM THE USE OF THIS PRODUCT ON PLANT SPECIES NOT RECOMMENDED ON THIS LABEL.

This product may be used on the following established plant species when container grown or field grown:

This product may be used on the following established plant species when container grown or field grown:							
TREES							
Balsam Fir White Fir Albus Flowering Maple Luteus Flowering Maple Tangerine Flowering Maple Tangerine Flowering Maple Vesuvius Red-Flowering Maple Flame Maple Norway Maple Red Maple Red Sunset Maple Silver Maple Sugar Maple Queen Palm River Birch Paper Birch	European White Birch Bottle Tree Black Olive Chinese Chestnut Carob Redbud Filicoides-Fernspray Cypress Gracilis-Slender Hinoki Cypress Swara False Cypress Squarrosa-Moss Cypress Palm (<i>Chamaedorea castarctarum</i>) Palm (<i>Chamaedorea costaricana</i>) Parlor Palm Cloud Nine Dogwood Flowering Dogwood	Kousa Dogwood Green Hawthorn Carrot Wood Arizona Cypress Russian Olive Redgum Eucalyptus Mealy Eucalyptus Silver Dollar Eucalyptus Coolibah Tree Red Ironbark Eucalyptus Ficus Mini Fucus White Ash Shamel Ash Ginko-Maidenhair Tree	HILES Honey Locust Shademaster Honey Locust Toyon Florida Anise-Tree Eastern Red Cedar Japanese Larch American Sweet Gum Tuliptree Southern Magnolia Crabapple White Mulberry Banana Blackgum Sourwood Norway Spruce	Pendula-Weeping Norway Spruce Repens-Spreading Norway Spruce Dwarf Alberta Spruce White Spruce Glauca-Colorado Blue Spruce Hoopsii-Hoop's Blue Spruce Hoopsii-Hoop's Blue Spruce Bristlecone Pine Canary Island Pine Shore Pine, Beach Pine Eldarica Pine Bosnian Pine Pumilio-Shrubby Swiss Mountain Pine Austrian Black Pine	Monterey Pine Red Pine White Pine Columnar Scotch Pine Scotch Pine Loblolly Pine Japanese Black Pine London Planetree American Sycamore California Sycamore Podocarpus Cottonwood Chilean Mesquite Yoshino Flowering Cherry Douglasfir	Scarlet Oak Bear Oak Pin Oak Willow Oak Red Oak Live Oak Black Locust Willow Giant Sequoia Mahogany Yellow Tab Baldcypress Eastern Hemlock Chinese Elm Mexican Fan Palm	
			ORNAMENTAL SHRU	IBS			
Edward Goucher Abelia Glossy Abelia Abyssinica Acacia Prostrate Acacia Shoestring Acacia Copper Leaf Amur Maple Coral Bark Japanese Maple Dwarf Japanese Maple Dwarf Japanese Maple Century Plant False Spirea Japanese Painted Fern Coyotebush William Penn Barberry Mentor Barberry Mentor Barberry Atropurea-Redleaf Japanese Barberry Aurea-Golden Japanese Barberry Aurea-Golden Japanese Barberry Crimson Pygmy Barberry Rose Glow Barberry Barbara Karst California Gold Pink Pixie Scarlet O'Hara Temple Fire Texas Dawn	Japanese Boxwood Korean Boxwood Common Boxwood Lemon Botwood Spring Torch Scotch Heather Japanese Camellia Sasanqua Camellia Feathery Cassia Wild Lilac Plum Yew Snow-in-Summer Kosteri Cypress Nana-Dwarf Hinoki Cypress Torulosa Cypress Filifera-Thread Cypress Areca Palm Summersweet Japanese Cleyera Sibirica-Siberian Dogwood Baileyi Red Osier Dogwood Flaviramea-Yellowtwig Dogwood Royal Purple Smoke Tree Eichholz Smoke Tree	Praecox-Early Cotoneaster Cranberry Cotoneaster Pyrenees Cotoneaster Himalayan Cotoneaster Rock Cotoneaster Zabel Cotoneaster Sago Palm Hollandia-Warminster Broom Lena-Scotch Broom Fragrant Daphne Deutzia Hopseed Bush Fruitland Silver Berry Purple Bell Heather Mediterranean Pink Heather Cornish Heather Winged Euonymus Canadale Gold Euonymus Emerald'n Gold Euonymus Sunspot Euonymus Wintercreeper Euonymus Variegated Evergreen Euonymus Variegated Evergreen Euonymus Spreading Euonymus	Pineapple Guava Forsythia August Beauty Gardenia Gardenia Radican Gardenia Salal/Lemon Leaf Carolina Jessamine Woadwaxen Ross Estey-Hibiscus Rose-of-Sharon- Heart Rose-of-Sharon- Heart Rose-of-Sharon- Head Bird Rose-of-Sharon- Woodbridge Holly Mystery Gardenia Henry Garnet Holly Leaf Sweetspire Ixora Juniper Mountain Laurel Crape Myrtle Lantana Coast Leucothoe Drooping Leucothoe Privet Chinese Fountain Palm Flowering Woodbine	Serotina Woodbine Trumpet Honeysuckle Leather Leaf Mahonia Creeping Mahonia Wax Myrtle Compacta-Dwarf Heavenly Bamboo Harbour Dwarf-Heavenly Bamboo Heavenly Bamboo Nana Compacta-Heavenly Bamboo Nana Compacta-Heavenly Bamboo Nana Compacta-Heavenly Bamboo Moato Dwarf Heavenly Bamboo Nana Purpurea-Heavenly Bamboo Nana Purpurea-Heavenly Bamboo Moods Dwarf Heavenly Bamboo Hardy Red Oleander Oleander Ruby Lace Oleander Fortunes Osmanthus Mockorange Pigmy Date Palm Fraser's Photinia Japanese Andromeda Mountain Fire Lily-of-the-Valley Snowdrift Lily-of-the-Valley Templebells Lily-of-the-Valley Valley Valentine Lily-of-the-Valley Forest Flame Lily-of-the-Valley	Mugo-Mugho Pine Green Pittosporum Japanese Pittosporum Wheeler's Dwarf Pittosporum Blue Cape Plumbago Plumbago Yewpine Tassel Fern Cinquefoil Carolina Laurel Cherry Dwarf Pink Flowering Almond Pyracantha Charisma-Monruce Rhaphiolepis Enchantress-Moness Rhaphiolepis India Hawthorn Springtime-Monme Rhaphiolepis Roundleaf Rhaphiolepis Azalea/Rhododendron Africa Sumac Ramanas Rose Rosemary Japanese Skimmia Reeve's Skimmia Anthony Waterer Spiraea Dolchia Spiraea	Japanese Alpine Spiraea Shirobana Spiraea Bridal Wreath Chinese Lilac Common Lilac Upright Japanese Yew Anglojap Yew Cape Honeysuckle <i>Ternstroemia gymnanthera</i> American Arborvitae Globosa-Globe Arborvitae Little Giant-Dwarf Arborvitae Nigra-Dark American Arborvitae Nigra-Dark American Arborvitae Pyramidalis-Pyramid Arborvitae Rheingold Arborvitae Aureus Nana-Dwarf Golden Arborvitae Minima Glauca-Dwarf Golden Arborvitae Christmas Palm Virburnum Weigela Xylosma Yucca	
			GROUND COVERS	3			
Wooly Yarrow Lily-of-the-Nile Beechgrass Cape Weed Thrift Asparagus Fern Bellflower Variegated Carex	Largeleaf loeplant Dwarf Plumbago Rockrose Coreopsis Crown Vetch Pampas Grass Cotoneaster White Iceplant	Descampsia Trailing Rosea Iceplant Iceplant Blue Fescue Strawberry, Beach Gazania Golden Hakonechloa Algerian Ivy	English Ivy Daylilly Rupturewort Albo-Marginata Hosta Aaronsbeard St. Johnswort Angelwing Jasmine Trailing Iceplant	White Lily Turf Big Blue Lily Turf Lilac Beauty Lily Turf Magestic Lily Turf Silvery Sunpoof Lily Turf Variegated Lirope Lily Turf Green/Creeping Lily Turf Eulalia Grass	Creeping Wirevine Myoporum Dwarf Mondo Grass Mondo Grass Trailing African Daisy Japanese Spurge Fountain Grass	Ribbon Grass Stonecrop (Sedum) Germander Asaian Jasmine Verbena Speedwell Periwinkle	
			ESTABLISHED FLOW	-			
Yarrow Floss Flower Alyssum Snapdragon African Daisy Dusty Miller Aster (perennial) Calendula	Comflower Velvet Centaurea Sweet Sultan Chrysanthemum Morningglory Coreopsis Cosmos Dahlia	Dianthus Marigold, Cape Snow on the Mountain Gaillardia Geum Gladiolus Baby's Breath	Sunflower Balsam Impatiens Ixora Sweet Pea Statice Lobelia ORNAMENTAL BUL		Petunia Phlox Portulaca Rose Blackeyed Susan Golden Glow Salvia	Pincushion Flower Lamb's Ears Stoke's Aster Marigold Nasturtium Vinca Zinnia	
This product may be applied for control of susceptible annual weeds in ornamental bulbs, e.g., bulbous iris, daffodil (narcissus), hyacinth and tulip. Apply to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. This product may also be applied following bulb emergence. For fall planted bulbs, apply this product again in later winter or early spring to weed-free soil surfaces.							

NON-BEARING FRUIT AND NUT TREES AND NON-BEARING VINEYARDS IN ORNAMENTAL LANDSCAPING ONLY*

Almond	Boysenberry	Fig	Kiwi	Orange	Pomegranate
Apple	Cherry, sour	Filbert	Lemon	Peach	Prune
Apricot	Cherry, sweet	Gooseberry	Loganberry	Pear	Raspberry
Avocado	Currant	Grape, American	Macadamia Nut	Pecan	Walnut, black
Blackberry	Dewberry	Grape, European	Nectarine	Pistachio	Walnut, English
Blueberry	Elderberry	Grapefruit	Olive	Plum	•

*Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants which will not bear fruit for at least one year after treatment.

NON-CROPLAND AREAS

This product is recommended as a preemergence herbicide for control of certain annual grasses and broadleaf weeds on industrial sites, utility substations, highway guard rails, sign posts and delineators. Apply anytime prior to germination of target weeds. Areas to be treated should be free of established weeds or existing weeds should be controlled with postemergence herbicides. Refer to the "General Information" section prior to using this product on non-cropland.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store product in its original container in a cool, dry, locked place out of the reach of children and animals

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill ration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The Manufacturer warrants that this product conforms to the chemical description of the label and is reasonably fit for the purpose stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. THE MANUFACTURER MAKES NO OTHER EXPRESSED OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESSED OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Manufacturer or the Seller. All such risks shall be assumed by Buyer.

WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions not reasonably foreseeable to Manufacturer, and Buver assumes the risk of any such use. Manufacturer, and Buyer assumes the risk of any such use.





LEBANON SEABOARD CORPORATION 1600 EAST CUMBERLAND STREET LEBANON, PA 17042