

LABEL SCAN CONTROL SHEET

5-14
2

Label ID 6798

Accepted Date (mm / dd / yyyy) 03 / 20 / 2001

EPA # 10404-82

Product Name LESCO PRE-M PLUS 0.86% PENDIMETHALIN VAR. FERT. COMBINATIONS

Product ID 27342

Reg Status Code R

Reg Status Text RENEWED

Use Code L

Use Text TURF AREA

Restriction Code

Restriction Text

Payor Name LESCO, INC.

Payor ID 10404

File Name LABEL 6798

Comments:

Scanning Action:

Completed OK

Rejected # Rejection

PRE-M[®] 0.86% Plus Fertilizer

ACCEPTED
VIA NOTIFICATION
EFFECTIVE DATE NOT REVIEWED

MAR 20 2001

PREEMERGENT WEED CONTROL

For Use on Turfgrasses

Not for use on turf being grown for sale or for other commercial use as sod, or for commercial seed production, or for research purposes.

- Preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed, and *Poa annua*.
- Provides quick green-up from soluble nitrogen.
- Prevents annual grassy and broadleaf weeds.

Active Ingredient:

Pendimethalin, N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzamine*0.86%

Inert Ingredients: 99.14%

Total 100.00%

*Licensed under one or more of the following U.S. Patents: 3,920,742; 4,066,441.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See below for additional precautionary statements

In case of emergency endangering life or property involving this product, call day or night, Chem-Trec 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water by cleaning of equipment or disposal of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at time of pesticide application.

Observe all cautions and limitations in this leaflet and on the labels of products used in combination with PRE-M 0.86% Plus Fertilizer herbicide. Do not use PRE-M 0.86% Plus Fertilizer herbicide other than in accordance with the instructions set forth on this label. The use of PRE-M 0.86% Plus Fertilizer herbicide not consistent with this label can result in injury to turf, animals, or persons. Keep container closed to avoid spills and contamination. DO NOT apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Storage: DO NOT store above 120°F. Store in original containers and keep closed; reseal opened bag by folding top down and securing. Store in a cool, dry place.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

PRE-M 0.86% Plus Fertilizer is a combination preemergent herbicide and fertilizer formulated to provide control of most annual grasses and certain broadleaf weeds found in turfgrass. PRE-M 0.86% Plus Fertilizer is labeled for use on the following turfgrass species: Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass, *Poa annua* (except putting green and tees), bermudagrass, St. Augustinegrass, bahiagrass, centipedegrass, and zoysiagrass.

PRE-M 0.86% Plus Fertilizer controls weeds as they germinate but will not control established weeds.

Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

Applied according to label directions and under normal growing conditions, PRE-M 0.86% Plus Fertilizer will not cause turf injury. Over-application can result in turf stand loss, turf injury, or soil residues. Uneven application can decrease weed control or cause turf injury.

Seeding diseases, cold weather, excessive moisture, high soil pH, high soil salt concentration, or drought can weaken turfgrass and increase the possibility of turf damage from PRE-M 0.86% Plus Fertilizer.

APPLICATION

Apply product uniformly at the recommended rate using a properly calibrated spreader (see table below for approximate settings). The efficacy of PRE-M 0.86% Plus Fertilizer will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. This will remove the product from the turfgrass foliage and establish a preemergent herbicide barrier in the soil. Avoid contact of product with porous surfaces such as driveways, stone, or wood. Removing particles will help prevent staining.

X-X-X

GUARANTEED ANALYSIS

Total Nitrogen (N)..... XX.XX%

X.XX% Ammoniacal Nitrogen

XX.XX% Urea Nitrogen*

Available Phosphate (P₂O₅).....X.XX%

Soluble Potash (K₂O).....XX.XX%

DERIVED FROM: xxxxxx, xxxxxxx, xxxxxx

Chlorine (Cl) Max. X.X%

*X.XX% Slowly available Nitrogen from LESCO® Poly Plus® Sulfur Coated Urea

Made in U.S.A.

Produced under the following U. S. Patent: 5,750,130

F335

Net Weight: 50 lb (22.7 kg)

EPA Reg. No. 10404-82

EPA Est. No. 10404-OH-4 (M), 10404-FL-1 (S), 59197-CA-1 (C)

Letter in lot number indicates manufacturing site.

F1406

Manufactured by: LESCO, Inc.

15885 Sprague Road

Strongsville, Ohio 44136-1799

758073xxxxxx

#XXXXXX

SUGGESTED SPREADER SETTINGS

Spreader	Ground Speed	Recommended Rate per 1000 sq ft
LESCO Rotary	3 mph	5.3"
LESCO Calibration Gauge		
Cyclone or Spyrker		

* Delivers 1.5 lb a/A and X.XX lb N/1,000 sq ft. ** Delivers 2.0 lb a/A and X.XX lb N/1,000 sq ft.

IMPORTANT: Spreader settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation in application rate. Spreader should be calibrated prior to each application by the operator.

Turfgrass Species Weeds Rates of PRE-M 0.86% Plus Fertilizer Comments:

Turfgrass Species	Weeds	Rates of PRE-M 0.86% Plus Fertilizer	Comments:
NORTHERN GRASSES			
Kentucky bluegrass	crabgrass foxtail	4.0 to 5.3 lbs/1,000 sq ft as the initial application prior to weed germination in the spring.	Apply a repeat application of 2.7 to 4.0 lbs/1,000 sq ft after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
Perennial ryegrass	<i>Poa annua</i> barnyardgrass		
Fine fescue	fall panicum oxalis		
Tall fescue	prostrate spurge purslane knotweed evening primrose hop clover		
	goosegrass	4.0 to 5.3 lbs/1,000 sq ft in the spring prior to weed germination.	Apply a repeat application of 4.0 lbs/1,000 sq ft

	crabgrass foxtail	4.0 lbs/1,000 sq ft in the spring prior to weed germination.	Apply a repeat application of 2.7 to 4.0 lbs/1,000 sq ft after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	<i>Poa annua</i> barnyardgrass		
	fall panicum oxalis		
	prostrate spurge purslane knotweed evening primrose hop clover		
	goosegrass	4.0 lbs/1,000 sq ft prior to weed germination in the spring.	Apply a repeat application of 4.0 lbs/1,000 sq ft after a minimum of 4 weeks.

	crabgrass foxtail	4.0 lbs/1,000 sq ft in the spring prior to weed germination.	Apply a repeat application of 2.7 to 4.0 lbs/1,000 sq ft after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	<i>Poa annua</i> barnyardgrass		
	fall panicum oxalis		
	prostrate spurge purslane knotweed evening primrose hop clover		
	goosegrass	4.0 lbs/1,000 sq ft prior to weed germination in the spring.	Apply a repeat application of 4.0 lbs/1,000 sq ft after a minimum of 4 weeks.

	crabgrass foxtail	4.0 lbs/1,000 sq ft in the spring prior to weed germination.	Apply a repeat application of 2.7 to 4.0 lbs/1,000 sq ft after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	<i>Poa annua</i> barnyardgrass		
	fall panicum oxalis		
	prostrate spurge purslane knotweed evening primrose hop clover		
	goosegrass	4.0 lbs/1,000 sq ft prior to weed germination in the spring.	Apply a repeat application of 4.0 lbs/1,000 sq ft after a minimum of 4 weeks.

	crabgrass foxtail	4.0 lbs/1,000 sq ft in the spring prior to weed germination.	Apply a repeat application of 2.7 to 4.0 lbs/1,000 sq ft after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	<i>Poa annua</i> barnyardgrass		
	fall panicum oxalis		
	prostrate spurge purslane knotweed evening primrose hop clover		
	goosegrass	4.0 lbs/1,000 sq ft prior to weed germination in the spring.	Apply a repeat application of 4.0 lbs/1,000 sq ft after a minimum of 4 weeks.

	crabgrass foxtail	4.0 lbs/1,000 sq ft in the spring prior to weed germination.	Apply a repeat application of 2.7 to 4.0 lbs/1,000 sq ft after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	<i>Poa annua</i> barnyardgrass		
	fall panicum oxalis		
	prostrate spurge purslane knotweed evening primrose hop clover		
	goosegrass	4.0 lbs/1,000 sq ft prior to weed germination in the spring.	Apply a repeat application of 4.0 lbs/1,000 sq ft after a minimum of 4 weeks.

	crabgrass foxtail	4.0 lbs/1,000 sq ft in the spring prior to weed germination.	Apply a repeat application of 2.7 to 4.0 lbs/1,000 sq ft after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	<i>Poa annua</i> barnyardgrass		
	fall panicum oxalis		
	prostrate spurge purslane knotweed evening primrose hop clover		
	goosegrass	4.0 lbs/1,000 sq ft prior to weed germination in the spring.	Apply a repeat application of 4.0 lbs/1,000 sq ft after a minimum of 4 weeks.

	crabgrass foxtail	4.0 lbs/1,000 sq ft in the spring prior to weed germination.	Apply a repeat application of 2.7 to 4.0 lbs/1,000 sq ft after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	<i>Poa annua</i> barnyardgrass		
	fall panicum oxalis		
	prostrate spurge purslane knotweed evening primrose hop clover		
	goosegrass	4.0 lbs/1,000 sq ft prior to weed germination in the spring.	Apply a repeat application of 4.0 lbs/1,000 sq ft after a minimum of 4 weeks.

	crabgrass foxtail	4.0 lbs/1,000 sq ft in the spring prior to weed germination.	Apply a repeat application of 2.7 to 4.0 lbs/1,000 sq ft after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	<i>Poa annua</i> barnyardgrass		
	fall panicum oxalis		
	prostrate spurge purslane knotweed evening primrose hop clover		
	goosegrass	4.0 lbs/1,000 sq ft prior to weed germination in the spring.	Apply a repeat application of 4.0 lbs/1,000 sq ft after a minimum of 4 weeks.

	crabgrass foxtail	4.0 lbs/1,000 sq ft in the spring prior to weed germination.	Apply a repeat application of 2.7 to 4.0 lbs/1,000 sq ft after a minimum of 4 weeks for extended control or where heavy weed infestations are expected.
	<i>Poa annua</i> barnyardgrass		
	fall panicum oxalis		
	prostrate spurge purslane knotweed evening primrose hop clover		
	goosegrass	4.0 lbs/1,000 sq ft prior to weed germination in the spring.	Apply a repeat application of 4.0 lbs/1,000 sq ft after a minimum of 4 weeks.

*Not for use on greens or tees

The efficacy of PRE-M 0.86% Plus Fertilizer will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. If PRE-M 0.86% Plus Fertilizer is not activated by rainfall or irrigation within 14 days, erratic weed control may result.

PRECAUTIONS

Use only on well established turfgrass with a dense and uniform stand. On newly planted areas, application should not be made until the turfgrass has filled in and has been mowed at least four times.

On turf that has been thinned or damaged due to winter injury, excessive moisture, etc., allow for turf recovery prior to making an application.

Applications made to overseeded warm-season turfgrasses may cause thinning or injury of the overseeded species.

Do not use on bentgrass or *Poa annua* (greens and tees) or injury may occur.

Delay reseeding or winter overseeding treated turfgrass for at least 2 months following application. Do not spring turfgrass for 3 months after application.

WARRANTY

LESCO warrants that this herbicide conforms to the chemical description on this label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn. RA Dept., 15885 Sprague Road, Strongsville, OH 44136-1799, referring to the item number found on this bag.

LESCO and Poly Plus are registered trademarks of LESCO, Inc.
G:\REGULWP\STRONGSV\Matr.LBPre-m086.snd.doc Rev. 11/20/00 WPT

(112298)